

## Influenza (FLU) Update for Week #50 Week Ending 12-15-18

**For week #50 (ending 12-15-18)** the CDC reported that **influenza (Flu)** activity, which includes diagnosed flu as well as **ILI (Influenza-Like-Illness)** increased and the threshold has been reached met to indicate that the 2018-19 Seasonal Flu season has begun. The percentage of respiratory specimens testing positive for flu more than quadrupled from last week to 11.0%.

Influenza A viruses are the most common virus found in sample testing, with Influenza A(H1N1)pdm09 the most common, except for Region 4 (the Southeast), which had A(H3N2) dominate. Influenza A(H2N2) and Influenza B viruses were considered to be co-circulating. Also, the majority of the flu viruses matched well with the 2018/19 Flu Vaccine.

See this link for more details on Flu and ILI activity, including charts, graphs and maps:  
<https://www.cdc.gov/flu/weekly/index.htm>

**FirstWatch RIN (Regional Influenza Network)** Alerts increased a small amount last week.

### **For the recently reported week ending December 15, 2018, the CDC reported:**

- Influenza-like illness (ILI) visits to clinics & other non-hospital facilities was 2.7% (l. w. 2.2%) & is above the national baseline of 2.2%. Eight (8) of the 10 regions (all except Regions 5 & 6) reported ILI at or above their region-specific baselines; it was 5 of 10 Regions last week. Two (2) states had high ILI activity; New York City, and nine (9) states had moderate ILI activity; Puerto Rico and 11 states reported low levels; and Washington DC and 28 states had minimal ILI activity.
- Flu cases indicating spread (documented by positive flu tests on respiratory specimens) were reported as Widespread in Guam and six (6) states; Regional in 18 states; 19 states reported local activity; Washington DC, Puerto Rico, the US Virgin Islands and seven (7) states reported sporadic flu activity. Clinical lab testing for influenza was positive for 11% of specimens, compared to 2.4% last week.
- Influenza A remained the dominant flu for 94.6% of the flu tests reported (92.2% last week), with 81.9% (80.9% last week) as A (H1N1)pdm09 viruses and the H3N2 subtype at 18.1% (19.1% last week). The rest of the tests showed 5.4% (7.8% l.w.) tested as Influenza B viruses, with 42.9% (66.7% l.w.) of Yamagata lineage and 57.1% (33.3% l.w.) Victoria lineage. This shows a very small increase in Influenza A viruses and therefore slightly less Influenza B. Typically, Influenza B viruses cause less severe illness and occur more towards the Spring.

All the flu viruses that were tested were susceptible to the antivirals oseltamivir, zanamivir, and peramivir (Tamiflu, Relenza, and Rapivab, respectively). There was high resistance to the adamantane group of antivirals, namely Amantadine and rimantadine (Flumadine), which are not effective for any Influenza B viruses). Baloxavir marboxil, under the brand name Xofluza, is a new influenza antiviral drug approved in October, 2018, as a new alternative. It also works on Influenza B viruses and controls the virus in a different way.

The CDC recommends as early as possible treatment with antivirals for those confirmed or suspected flu with severe, complicated, or progressive disease, who are hospitalized, or at high risk for complications of flu. See this link for a list of those at risk for complications from flu:

[https://www.cdc.gov/flu/about/disease/high\\_risk.htm](https://www.cdc.gov/flu/about/disease/high_risk.htm)

**The CDC provides an interactive U.S. map that will link to each state's public health authorities. ILI and Flu information and processes, as well as other diseases and public health topics. This site includes a tremendous amount of information at the State and even Local level.**

Find it at this site: <https://www.cdc.gov/flu/weekly/usmap.htm> (currently represents Week #50)

## -- For Influenza-Like Illness:

**High ILI Activity: (2 states):** Colorado and Georgia

**Moderate ILI Activity: (New York City & 9 states):** Alabama, Arizona, Arkansas, Kentucky, Louisiana, Missouri, New Jersey, South Carolina, and Virginia

**Low Activity: (Puerto Rico & 11 states):** California, Massachusetts, Minnesota, Mississippi, New Mexico, New York, North Carolina, Oklahoma, Pennsylvania, Texas, and Utah

**Minimal Activity: (Washington D.C., & 28 states):** Alaska, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Maryland, Michigan, Montana, Nebraska, Nevada, New Hampshire, North Dakota, Ohio, Oregon, Rhode Island, South Dakota, Tennessee, Vermont, Washington, West Virginia, Wisconsin, and Wyoming

## -- For Flu (positive flu tests):

**Widespread Activity: (Guam & 6 states):** Alabama, California, Delaware, Georgia, Massachusetts and New York

**Regional Activity (18 states):** Arizona, Connecticut, Florida, Idaho, Kentucky, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, Texas, Vermont, and Virginia

**Local Activity (19 states):** Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, North Dakota, Oklahoma, South Carolina, Tennessee, Utah, and Wyoming

**Sporadic Activity: (Washington D.C., Puerto Rico, U.S. Virgin Islands, & 7 states):** Alaska, Hawaii, Maine, South Dakota, Washington, West Virginia, and Wisconsin

## -- Other Data:

The Hospitalization rate from Flu was 2.9 per 100,000 (last week 1.9%). Children (ages 0-4) had the highest hospitalization rate of 7.7 per 100,000 (last week 5.0%), with older adults (age  $\geq 65$  years) at 6.6% per 100,000 (l.w. 4.6%); and adults (age 50-64 years) at 3.4% per 100,000 (l.w. 2.1%). Most, 84.0% (l.w. 78.9%), were caused by Influenza A viruses; 13.8% (l.w. 18.4%) were from Influenza B viruses; 1.3% (l.w. 1.8%) showed co-infection with both Influenza A and B viruses; and 1.0% (l.w. 0.9%) were not typed for a specific flu virus.

Death rates for pneumonia and influenza in adults (6.0%) is below the epidemic threshold of 6.7% for week #49. Death reports often aren't reported for data purposes the same week.

There was one pediatric death attributed to flu reported this week, for a total of 7 for this flu season

## -- Flu in Canada, Europe & the World:

### Canada:

According to the Public Health Agency of Canada (PHAC), throughout Canada there was a continued increase in flu activity for Week #50, week ending 12-15-18, with 22.8% of tests positive for flu. Influenza A is the predominant circulating Flu with 99% of specimens, and with A(H1N1)pdm09 as the dominant subtype (94%). Hospitalizations continue in all age ranges, but the highest estimated rate was for children less than 5 years old. A Flu Summary will not be posted for next week; a report on Jan 4, 2019 will include both Weeks # 51 & 52.

### **For more specific information see:**

**On this week's flu activity:** <https://www.canada.ca/en/public-health/services/publications/diseases-conditions/fluwatch/2018-2019/week50-december-9-december-15-2018.html>

### **Canadian Flu Information:**

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza.html>



Every Record. In Real Time. Automatically.

**General Page for Canadian Flu Watch Surveillance with links to different components:**

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance.html>

**About the Canadian Influenza Activity Surveillance System:**

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance/about-fluwatch.html>

### **Europe:**

According to the European Center for Disease Prevention & Control (ECDC), influenza activity was at baseline or low throughout the European Region, although most countries reported a geographic distribution of local, regional, or widespread flu. The majority of circulating virus remained Influenza A for patients both in hospital and out; 18% of tested specimens were positive for flu.

**For more information see:** <http://flunewseurope.org/>

### **World:**

The **World Health Organization (WHO)** provides info on Influenza in Member Countries **here:**

[https://www.who.int/influenza/surveillance\\_monitoring/en/](https://www.who.int/influenza/surveillance_monitoring/en/)

### **First Responder Specific Information**

There are many websites that may be helpful in planning and managing seasonal flu within First Responder organizations. A few of those websites are included here:

**NIOSH on Flu for Employers/Employees:**

<https://www.cdc.gov/niosh/topics/flu/>

**Protection from Flu:**

<https://www.cdc.gov/flu/protect/habits/index.htm>

**Weekly Flu Map:**

<https://www.cdc.gov/flu/weekly/usmap.htm>

**World Map Showing Flu & Other Infectious Diseases:**

<http://www.healthmap.org/en/>

### **Other Actions First Responders Should Consider**

- First Responders should be vaccinated for Flu each season to prevent getting flu themselves, taking it home to family members, or transmitting it to patients in their care. Family members and patients may be at increased risk of complications from flu.
- Perform proper hand hygiene including frequent handwashing and the use of hand sanitizers in general, and particularly when providing patient care or after touching surfaces.
- Masks (N95 or N100) should be used in the presence of patients with cough and/or fever; preferably before being within 6 feet of the patient. This becomes even more important if droplet producing procedures are being performed (i.e. suctioning, nebulizer treatments, BVM, intubation).
- Care should be taken to avoid touching your own face and mucous membranes (eyes, mouth, nose) since the flu virus is frequently found on surfaces such as door knobs, writing & recording tools (pens and tablets), cot and equipment handles, phones, light switches, as well as clothing, bed clothes, etc.
- Report signs/symptoms of flu to your physician or other appropriate provider for early assessment and care. Alert your employer per policy.

- Cough and sneeze into your sleeve, if a tissue is not available, and not onto your hands. Watch this Youtube video for a humorous but educational approach on the subject.  
<https://www.youtube.com/watch?v=CtnEwvUWDo0>
- Stay away from others if you are sick.
- Be aware of your exposure risk and history to prevent exposing others. Take extra precautions or avoid those with immunocompromise, when possible, if you have a known or likely exposure.
- Antivirals may be indicated for the treatment of flu, particularly for those in high risk groups, those who are hospitalized or have severe, complicated or progressing flu. Those that present with 48 hours of the onset of symptoms may also be given antivirals, based on PCP judgement but make sure the practitioner is aware of their First Responder Role. See <https://www.cdc.gov/flu/antivirals/whatyoushould.htm>

**And, for consideration when looking at yourself, your family and friends, or your patients, consider the following information regarding complications of flu:**

Flu is much more worrisome for the very young and the elderly, as well as those who fit into one of the high risk categories see this link for the list: [https://www.cdc.gov/flu/about/disease/high\\_risk.htm](https://www.cdc.gov/flu/about/disease/high_risk.htm). Signs of ILI/Flu in this group requires careful assessment to rule out complications and these groups are much more likely to need medical oversight to assure adequate care. Young children and those over 65 are typically at greater risk for complications, hospitalization, and even death.

Consideration should be given to perhaps monitoring these groups more closely, with inclination for more comprehensive assessment and transport for further evaluation, when presented with possible flu and any signs of complications.

Complications of flu, sometimes requiring hospitalization and even leading to death, tend to occur after the person has begun to get better from the flu and then appears to relapse. EMS personnel may want to look more closely at those patients when the call is not about the initial signs and symptoms of flu, but about increasing or different signs that have appeared, often from five days to two weeks after the initial flu symptoms began.

A study was published by the Institute for Clinical Evaluative Sciences in *NEJM* (*New England Journal of Medicine*). See details below:

## Flu infection may raise risk of heart attack, particularly in first 7 days



**Study confirms importance of flu vaccination for people at risk of heart disease.**

Researchers looked at nearly 20,000 Ontario adult cases of lab-confirmed influenza (2009-2014) and then identified 332 patients who were hospitalized for a heart attack within one year of flu diagnosis.



For this population, the risk of heart attack was **6 times higher** within the first week of a flu diagnosis.

Factors that may be associated with more risk:

- being age 65 and older
- infection with influenza B
- no previous heart attack

**The researchers say that people at risk of heart disease should take care to prevent flu through measures including handwashing and vaccination, and should not delay medical evaluation for heart symptoms, particularly in the first week of an acute respiratory infection.**

Kwong JC et al. *NEJM*. 2018.

**Institute for Clinical Evaluative Sciences**  
ices.on.ca

Public Health Ontario | Santé publique Ontario

Image courtesy of ICES/PHO



Every Record. In Real Time. Automatically.

“The researchers add that patients should not delay medical evaluation for heart symptoms particularly within the first week of an acute respiratory infection.” (Lisa Schnirring, News Editor: *CIDRAP News* ;Jan 25, 2018)  
For more information on Influenza and the Heart Attack Study, please see the link below.  
[https://www.eurekalert.org/pub\\_releases/2018-01/pho-rci011818.php](https://www.eurekalert.org/pub_releases/2018-01/pho-rci011818.php)

This document is provided by FirstWatch as a service to prehospital and public health providers. The information has been compiled from a variety of sources such as the CDC, Health Canada and WHO. It is copyrighted, but may be freely copied and distributed as long as it remains intact. For up-to-date, EMS specific information on emerging diseases, please visit [www.firstwatch.net/hi](http://www.firstwatch.net/hi)