

Influenza (FLU) Update for Week #47 -- Week Ending 11-23-19

The CDC reported that seasonal flu activity continued to increase over the previous week but the level of activity and the predominant influenza virus differ according to region. From a national perspective, ILI has been at/above baseline for three weeks. Some areas, particularly in parts of the South and West are seeing elevated levels, while other areas still have low levels of flu activity. According to the data, the 2019/20 flu season is ramping up earlier compared with other recent years (see chart below). The one recent exception is the 2009/10 season, displayed on the chart with a silver line, which represents the pandemic flu year. **Influenza-like illness (ILI) is defined as a temperature at/above 100° F [37.8° C] and cough and/or sore throat without a known cause other than flu. A Flu case, that is included in CDC data, indicates a positive flu test read by either a Clinical or Public Health Lab.**




Chart copied from the CDC URL <https://www.cdc.gov/flu/weekly/index.htm> ; content source Center for Disease Control and Prevention, National Center for Immunization and Respiratory Diseases (NCIRD); page last reviewed 12/2/19; chart accessed 12/2/19.

Diagnosed Flu and ILI are expected to increase over the next few weeks and months. **It is recommended by public health officials & health care providers that for all those 6 months or older, flu vaccinations should be completed ASAP unless they have a**

valid medical contraindication. Most will receive a vaccination that covers 4 strains of influenza.

The severity of Flu & ILI for this season, as well as which viruses will ultimately dominate, is still too early to identify, but there are hints that this season may be more severe than many, since most recent seasons have hit the national threshold for ILI typically between weeks #48 and #50, but this season hit the threshold at week #45 and has continued to climb. As more data is collected by the CDC further into the season, its severity and other report components will be included in this Summary.

See this link for more details, including charts, graphs, and maps:
<https://www.cdc.gov/flu/weekly/index.htm>

FirstWatch RIN (Reginal Influenza Network) Alerts increased over the previous week.

For the most recently reported week ending November 23, 2019, the CDC reported:

--ILI visits to clinics and other non-hospital facilities was at 2.9%, higher than last week's 2.5% and above the national baseline of 2.4%. The regional range was between 1.5% and 6.8% for Week #47. Four (4) of 10 Regions reported that outpatient visits for ILI were at/above their own regional baselines including **Region 3** (DC, DE, MD, PA, VA, & WV), **Region 4** (AL, FL, GA, KY, MS, NC, SC, & TN), **Region 6** (AR, LA, NM, OK & TX), and **Region 10** (AK, ID, OR & WA).

--Flu cases, documented by positive flu tests from Clinical Labs, showed that 8.0% of respiratory specimens tested for influenza were positive, which is higher than last week's report of 7.3%.

-- For this reporting period, viewed from a national standpoint, Influenza B/Victoria viruses were reported more frequently, than Influenza A(H1N1)pdm09 and Influenza A(H3N2) viruses, but are considered to be cocirculating. The predominant strain varied by region and patient age. To access specific state and regional information on circulating flu viruses, please see: <https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>

--Clinical Labs reported that of 21,367 specimens tested, 1,702 were positive for flu at 8.0% (last week 7.3%), with 70.4% Influenza B and 29.6% Influenza A. Public Health Labs reported data which has a more limited number of specimens (1,193 tested with 412 positive for flu) but provides a more specific breakdown of specific flu virus strains. Influenza B was detected at 60.4% and Influenza A at 39.6%. **Further breakdown revealed that Influenza B showed 98.8% Victoria lineage and 1.2% Yamagata, while H1N1 is the dominant A with 75.9% and H3N2 at 24.1%.** Influenza B viruses are typically seen in greater numbers nearer the end of the flu season rather than the beginning and may be a tougher flu for children.

--Vaccine Coverage: Testing of Influenza A(H1N1) samples revealed that all samples tested were well matched with the components in the 2019-20 flu vaccine, while 87.5% of A(H3N2) Influenza virus samples matched; 63.6% of B/Victoria virus samples were similarly matched. Testing of B/Yamagata samples were pending at the time of the CDC's report.

The CDC provides an interactive U.S. map that will link to each state's public health authorities, ILI and Flu information and processes, as well as other diseases and public health topics. This site includes a tremendous amount of information at the State, and even Local, level.

Find it at this site: <https://www.cdc.gov/flu/weekly/usmap.htm>

--For Influenza-Like Illness (ILI)

High ILI Activity (Puerto Rico & 7 state): Alabama, Georgia, Louisiana, Mississippi, South Carolina, Tennessee and Texas

Moderate ILI Activity (7 states): Arkansas, Connecticut, Nebraska, Nevada, New Mexico, Virginia, and Washington

Low Activity (New York City & 8 states): Arizona, Colorado, Hawaii, Maryland, Minnesota, New Jersey, Oklahoma, and Oregon

Minimal Activity (& 28 states): Alaska, California, Delaware, Florida, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Massachusetts, Michigan, Missouri, Montana, New Hampshire, New York, North Carolina, North Dakota, Ohio, Pennsylvania, Rhode Island, South Dakota, Utah, Vermont, West Virginia, Wisconsin, and Wyoming

Insufficient Data to Calculate: U.S. Virgin Islands & Washington D.C.

--For Flu (positive Flu tests)

Widespread Activity (10 states): Alabama, Alaska, California, Louisiana, Massachusetts, Nevada, New Mexico, South Carolina, Tennessee, and Texas

Regional Activity (14 states): Arizona, Colorado, Connecticut, Florida, Georgia, Idaho, Indiana, Kentucky, Nebraska, New York, Oregon, Pennsylvania, Virginia, and Washington

Local Activity (Puerto Rico & 19 states): Arkansas, Delaware, Illinois, Iowa, Maryland, Michigan, Minnesota, Mississippi, Montana, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, South Dakota, Utah, Vermont, and Wisconsin

Sporadic Activity (U.S. Virgin Islands, & 7 states): Hawaii, Kansas, Maine, Missouri, Rhode Island, West Virginia, and Wyoming

No Activity: (0 states):

No Report: Washington D.C. and Guam

--Other Data:

Hospitalizations from Flu: The hospitalization rate for all ages remained 2.0 per 100,000 and is typical for this time period when compared with previous seasons. The highest rate was in those aged ≥ 65 years (5.1/100,000), then children ages 0-4 (3.1/100,000), and then adults ages 50-64 (2.0/100,000). The vast majority were from infection with influenza A (62.6%) with an even split between H1N1(pdm09) and H3N2.

Death rates for pneumonia & influenza in adults: from data as of 11/27/19, 5.1% of adult deaths during week #46 (ending 11/16) were associated with influenza and pneumonia. This is below the epidemic threshold of 6.3% for that week. This data reporting typically lags a week or more behind the reporting week.

One (1) pediatric death associated with influenza was reported during week #47, making a total of five (5) flu-associated pediatric deaths for the 2019-20 Flu Season.

International:

Canada:

Flu in Canada for Week #47 (11/17 - 11/23, 2019):

According to the Public Health Agency of Canada (PHAC), influenza activity increased and is nationally now at the seasonal threshold level, indicating that the flu season started during Week #47, which is a typical start of the flu season. Influenza A(H3N2) continued to be the most common flu virus circulating, but the amount of A(H1N1) and Influenza B increased proportionally. Influenza B accounted for 1/3 of positive specimens, which is atypical for this early in the season

Localized Activity: in 4 Regions: Quebec (1), Ontario (1), Saskatchewan (1), Alberta (1)

Sporadic Activity: in 25 Regions: Newfoundland & Labrador (2), Prince Edward Island (1), New Brunswick (3), Quebec (4), Ontario (4), Manitoba (1), Saskatchewan (1), Alberta (4), British Columbia (4), Northwest Territory (1)

No Activity: in 20 Regions: Newfoundland & Labrador (2), Nova Scotia (4), New Brunswick (4), Quebec (1), Ontario (1), Manitoba (4), Saskatchewan (1), British Columbia (1), Yukon Territory (1) Northwest Territory (1)

No Data Reported: Ontario (1), Nunavut (3)

PHAC Flu Watch Summary & Influenza/ILI Activity Map for Week #47 see:

<https://www.canada.ca/en/public-health/services/publications/diseases-conditions/fluwatch/2019-2020/week-47-november-17-23-2019.html#a1>

PHAC Home Page for Surveillance on Flu: <http://www.phac-aspc.gc.ca/fluwatch/>

Public Health Agency of Canada (PHAC) Home Page for Information on Flu:

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza.html>

PHAC Provides Flu Watch Summary & Link to Full Influenza Reports Past & Present:

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance/weekly-influenza-reports.html>



Every Record. In Real Time. Automatically.

PHAC Interactive Map for Flu Activity Across Canada:

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance/map-activity-levels.html>

PHAC Site that Explains How to Become a Canadian Flu Watcher:

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/fluwatcher.html>

Europe:

European Center for Disease Prevention & Control - Flu for Week #47 (11/18 – 11/24, 2019):

Influenza activity was low or at baseline levels throughout Europe. 11.3% tested positive for flu from those going to sentinel PCPs for ILI and ARI. Flu viruses A & B, were found in specimens from those with respiratory illness seeking medical care, with more A than B, but with influenza B increasing. Data from 22 countries/regions reported mortality from all causes were at expected levels for this time of year.

For more info see: <http://flunewseurope.org/>

Public Health England Flu Summary:

<https://www.gov.uk/government/statistics/weekly-national-flu-reports>

Global:

World Health Organization (WHO) Weekly Flu Summary (FluNet):

http://www.who.int/influenza/gisrs_laboratory/flunet/en/

WHO Global Flu Overview with Map and Text:

https://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/

Global Epidemiology Reports:

WHO Collaborating Centers for Flu in [Australia](#), [Japan](#), & the [United Kingdom](#)

First Responder-Specific Information

There are many websites that may be helpful in planning and managing seasonal flu within First Responder organizations. A few of those websites are included here:

NIOSH on Flu for Employers/Employees:

<https://www.cdc.gov/niosh/topics/flu/>

NIOSH Listing on Absenteeism in the Workplace throughout the US:

<https://www.cdc.gov/niosh/topics/absences/default.html>

Protection from Flu:

<https://www.cdc.gov/flu/protect/habits/index.htm>

World Map Showing Flu & Other Infectious Diseases:

<http://www.healthmap.org/en/>

- First Responders should be vaccinated for Flu each season to prevent getting flu themselves, taking it home to family members or transmitting it to patients in their care. Family members and patients may be at increased risk of complications from flu.
- Perform proper hand hygiene including frequent handwashing with soap and water, and the use of hand sanitizers in general, and particularly when providing patient care or after touching surfaces.
- Masks (N95 or 100) should be used in the presence of patients with cough and/or fever. Procedure or surgical masks can also be used but may not provide the same level of protection.
- Care should be taken to avoid touching their own face and mucous membranes (eyes, mouth, nose) since the flu virus is frequently found on surfaces such as door knobs, cot and equipment handles, phones, as well as clothing, bed clothes, etc. Once picked up on the hands from touching inanimate objects or from the secretions of a patient, it can transmit the flu (or any respiratory infection virus) via the mucous membranes of the face and head.
- Report signs/symptoms of flu to your physician or other appropriate provider for early assessment and care. Signs & symptoms suggest a period of contagiousness and out your patients and co-workers at risk.
- Cough and sneeze into your sleeve, if a tissue is not available, and not onto your hands.

- Stay away from others if you are sick.
- Be aware of your exposure risk and history. Take extra precautions or avoid those with immunocompromise, when possible, if you have a known or likely exposure.
- Antivirals may be indicated for the treatment of flu, particularly for those in high risk groups, those who are hospitalized or have severe, complicated or progressing flu. Those that present with 48 hours of the onset of symptoms may also be given antivirals, based on PCP judgement but make sure the practitioner is aware of their Frist Responder Role. See <https://www.cdc.gov/flu/antivirals/whatyoushould.htm>
- Flu is much more worrisome for the very young and the very old. Signs of ILI in this group requires careful assessment to rule out complications and these groups are much more likely to need to be transported to assure adequate care. Young children and those over 65 are typically at greater risk for complications, hospitalization, and even death. Hospitalization rates may also be elevated in those aged 50-64. Consideration should perhaps be given to monitoring these groups more closely, with inclination for more comprehensive assessment and transport for further evaluation, when presented with possible flu and any signs of complications.

Complications of flu, sometimes requiring hospitalization and even leading to death, tend to occur after the person has begun to get better from the flu and then appears to relapse. EMS personnel may want to look more closely at those patients when the call is not about the initial signs and symptoms of flu, but about increasing or different signs that have appeared.

- A study was published by the Institute for Clinical Evaluative Sciences in *NEJM*; see details below

Flu infection may raise risk of heart attack, particularly in first 7 days



Study confirms importance of flu vaccination for people at risk of heart disease.

Researchers looked at nearly 20,000 Ontario adult cases of lab-confirmed influenza (2009-2014) and then identified 332 patients who were hospitalized for a heart attack within one year of flu diagnosis.



For this population, the risk of heart attack was **6 times higher** within the first week of a flu diagnosis.

Factors that may be associated with more risk:

- being age 65 and older
- infection with influenza B
- no previous heart attack

The researchers say that people at risk of heart disease should take care to prevent flu through measures including handwashing and vaccination, and should not delay medical evaluation for heart symptoms, particularly in the first week of an acute respiratory infection.

Kwong JC et al. *NEJM*. 2018.

Institute for Clinical Evaluative Sciences

ices.on.ca

Public Health Ontario | Santé publique Ontario

Image courtesy of ICES/PHO “The researchers add that patients should not delay medical evaluation for heart symptoms particularly within the first week of an acute respiratory infection.” (Lisa Schirring, News Editor: *CIDRAP News*; Jan 25, 2018)

For more information on the Influenza and Heart Attack Study, please see the link below.
https://www.eurekalert.org/pub_releases/2018-01/pho-rci011818.php